


## SUPPLEMENTARY MATERIAL S1



**FIGURE S1** | Ontogenetic development of the lateral-line canals in larvae and juveniles of *Electrophorus varius*, in lateral and dorsal views. **A.** 1.9 cm TL; **B.** 2.5 cm TL; **C.** 3.0 cm TL; **D.** 5.0 cm TL; **E.** 9.0 cm TL; **F.** 38.0 cm TL. IO: infraorbital canal; MD: mandibular canal; O: otic canal; PO: preopercular canal; SO: supraorbital canal; ST: supratemporal canal. The red arrow indicates the position of the otic pit in the specimens.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2021 The Authors. Diversity and Distributions Published by SBI

- Verçoza G, Shibuya A, Bastos DA, Zuanon J, Rapp Py-Daniel LH. Organization of the cephalic lateral-line canals in *Electrophorus varius* de Santana, Wosiacki, Crampton, Sabaj, Dillman, Mendes-Júnior & Castro e Castro, 2019 (Gymnotiformes: Gymnotidae). Neotrop Ichthyol. 2021; 19(2):e200075. <https://doi.org/10.1590/1982-0224-2020-0075>