

SUPPLEMENTARY MATERIAL S5

TABLE S5 | Ranges in millimeters (mm) and proportions of disc width (%DW) of clasper and tail measurements taken from adults, subadults, juvenile and neonate specimens of *Paratrygon aiereba*; (N) corresponds to the number of specimens analyzed. Indicar os lotes e são tipos.

Clasper	Adults					Subadults				Juveniles				Neonates						
	mm		%DW		N	mm		%DW		N	mm		%DW		N	mm		%DW		N
External length of clasper	40	60	6.61	10.4	8	12	3.1	1	5	16	2.3	4.4	13	4	5	2.4	2.7	3		
Internal length of clasper	81	113	15.1	21.5	7	31	8.1	1	15	50	6.8	11.7	13	11	15	6.7	7.5	3		
Dorsal pseudosiphon length	9	12	1.5	2.0	8	–	–	0	1	4	0.4	0.9	3	–	–	–	–	0		
Ventral pseudosiphon length	24	52	4.7	7.6	7	–	–	0	7	15	2.1	3.5	2	–	–	–	–	0		
Tail																				
Tail width	32	55	6.0	7.3	12	18	27	5.9	7.0	2	12	32	5.0	7.7	24	11	15	5.5	8.8	7
Distance between cloaca and tail tip	115	545	19.5	87.5	11	182	60.1	1	101	341	35.5	159.2	23	82	331	41.0	183.0	7		
Distance between cloaca and caudal sting	103	191	19.4	26.4	10	53	17.5	1	44	100	19.6	26.6	23	42	60	21.0	31.3	7		



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons **CC-BY 4.0**

© 2021 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Loboda TS, Lasso CA, Rosa RS, Carvalho MR.** Two new species of freshwater stingrays of the genus *Paratrygon* (Chondrichthyes: Potamotrygonidae) from the Orinoco basin, with comments on the taxonomy of *Paratrygon aiereba*. Neotrop Ichthyol. 2021; 19(2):e200083. <https://doi.org/10.1590/1982-0224-2020-0083>



Official Journal of the Sociedade Brasileira de Ictiologia