

## SUPPLEMENTARY MATERIAL S5

**TABLE S5 |** D-loop based pairwise  $\Phi_{ST}$  index for *Pseudobatos percellens*. Sampled from the three states of Brazil, being Florianópolis/SC (SC), Pontal do Paraná/PR (PR), Cananéia/ SP (SP1), Mongaguá/SP (SP2), Santos/SP (SP3). The values in bold were significant ( $p$  value < 0.05).

|     | SC            | PR             | SP1           | SP2    | SP3 |
|-----|---------------|----------------|---------------|--------|-----|
| SC  | -             |                |               |        |     |
| PR  | <b>0.3650</b> | -              |               |        |     |
| SP1 | 0.1315        | <b>0.3760</b>  | -             |        |     |
| SP2 | 0.1990        | <b>0.2254</b>  | <b>0.1995</b> | -      |     |
| SP3 | 0.1616        | <b>s0.2255</b> | <b>0.1498</b> | 0.0320 | -   |

- Cruz VP, Adachi AMCL, Oliveira PH, Ribeiro GS, Paim FG, Souza BC, Rodrigues ASF, Vianna M, Delpiani SM, Díaz de Astarloa JM, Rotundo MM, Mendonça FF, Oliveira C, Lessa RP, Foresti F. Genetic diversity in two threatened species of guitarfish (Elasmobranchii: Rhinobatidae) from the Brazilian and Argentinian coasts: an alert for conservation. *Neotrop Ichthyol*. 2021; 19(2):e210012. <https://doi.org/10.1590/1982-0224-2021-0012>