

SUPPLEMENTARY MATERIAL S3

ALIMENTARY ACTIVITY

Stomach fullness (percent of fullness) was visually estimated on a scale from 0 to 3 (Hahn *et al.*, 1999) considering the amount of food filling the stomach: 0 (empty), 1 (< 25%), 2 (25–75%) and 3 (> 75%). We reported a high number of empty stomachs during the sampled period (total average – 76.58%). The Fig. S3 shows the number of each species. Acestrorhynchids species showed a higher number of empty stomachs in pre-damming, while cynodontids showed an increased number in post-damming.




FIGURE S3 | Stomach fullness of *Acestrorhynchus falcirostris*, *A. heterolepis*, *Hydrolycus scomberoides*, and *Rhaphiodon vulpinus* collected from Madeira River, during sampling periods before and after damming.



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2021 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Lonardoni AP, Röpkke CP, Melo T, Torrente-Vilara G.** Damming in the Madeira River modifies the food spectrum of piscivorous and affects their resource partitioning. *Neotrop Ichthyol.* 2021; 19(3):e210087. <https://doi.org/10.1590/1982-0224-2021-0087>