

## SUPPLEMENTARY MATERIAL S4

TABLE S4 | SIMPER analysis – Species that contribute most to the dissimilarities between seasons (Bray Curtis), in descending order.

Species	High-water average biomass	High-water average biomass	Average dissimilarity	% Contribution	Cumulative %
<i>Mylossoma albiscopum</i>	23.37	18.41	11.76	14.64	14.64
<i>Pimelodus blochii</i>	16.55	2.84	8.47	10.54	25.18
<i>Colossoma macropomum</i>	15.74	2.23	8.08	10.05	35.23
<i>Triportheus angulatus</i>	1.24	12.10	5.96	7.42	42.66
<i>Osteoglossum bicirrhosum</i>	3.57	10.48	5.52	6.87	49.53
<i>Pseudoplatystoma punctifer</i>	0.93	11.12	5.51	6.85	56.38
<i>Arapaima gigas</i>	1.23	9.82	5.18	6.45	62.83
<i>Phractocephalus hemiliopterus</i>	1.60	7.28	3.89	4.84	67.67
<i>Triportheus auritus</i>	2.52	7.46	3.88	4.83	72.50
<i>Pygocentrus nattereri</i>	7.81	0.68	3.70	4.60	77.11
<i>Pterygoplichthys</i> spp.	3.19	5.75	3.57	4.44	81.55
<i>Semaprochilodus insignis</i>	5.59	0.82	2.96	3.69	85.23
<i>Schizodon fasciatus</i>	4.45	2.24	2.58	3.21	88.44
<i>Prochilodus nigricans</i>	2.56	3.12	2.44	3.03	91.48



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2021 The Authors.  
Diversity and Distributions Published by SBI



Official Journal of the  
Sociedade Brasileira de Ictiologia

## HOW TO CITE THIS ARTICLE

- Tregidgo D, Parry L, Barlow J, Pompeu PS. Urban market amplifies strong species selectivity in Amazonian artisanal fisheries. *Neotrop Ichthyol.* 2021; 19(3):e210097. <https://doi.org/10.1590/1982-0224-2021-0097>