

SUPPLEMENTARY MATERIAL S3

References S3 | References used to obtain data for the classification of trophic groups and life-history (based on Brejão *et al.*, 2013 and Zeug, Winemiller, 2007) of fish from nine streams in the Amazon basin, in the state of Mato Grosso, Brazil.

- **Albert JS, Crampton WGR.** Diversity and phylogeny of Neotropical electric fishes (Gymnotiformes). In: Bullock TH, Hopkins CD, Popper AN, Fay RR, editors. *Electroreception*. New York: Springer; 2006. p.360–409. http://dx.doi.org/10.1007/0-387-28275-0_13
- **Arantes CC, Winemiller KO, Petrere M, Castello L, Hess LL, Freitas CEC.** Relationships between forest cover and fish diversity in the Amazon River floodplain. *J Appl Ecol*. 2017; 55(1):386–95. <https://doi.org/10.1111/1365-2664.12967>
- **Boersma KS, Dee LE, Miller SJ, Bogan MT, Lytle DA, Gitelman AI.** Linking multidimensional functional diversity to quantitative methods: A graphical hypothesis-evaluation framework. *Ecology*. 2016; 97(3):583–93. <https://doi.org/10.1890/15-0688>
- **Bojsen BH, Barriga R.** Effects of deforestation on fish community structure in Ecuadorian Amazon streams. *Freshwa Biol*. 2002; 47(11): 2246–60. <https://doi.org/10.1046/j.1365-2427.2002.00956.x>
- **Brejão GL, Gerhard P, Zuanon J.** Functional trophic composition of the ichthyofauna of forest streams in eastern Brazilian Amazon. *Neotrop Ichthyol*. 2013; 11(2):361–73. <http://dx.doi.org/10.1590/S1679-62252013005000006>
- **Burress ED.** Cichlid fishes as models of ecological diversification: Patterns, mechanisms, and consequences. *Hydrobiologia*. 2015; 748(1):7–27. <https://doi.org/10.1007/s10750-014-1960-z>
- **Carvalho LN, Zuanon J, Sazima I.** Natural history of Amazon fishes. In: Del-Claro K, Oliveira PS, Rico-Gray V, editors. *Tropical biology and conservation management: case studies*. Oxford: Eolss Publishers; 2009. Available from: <http://www.eolss.net>
- **Carvalho RA, Tejerina-Garro FL.** The influence of environmental variables on the functional structure of headwater stream fish assemblages: A study of two tropical basins in Central Brazil. *Neotrop Ichthyol*. 2015; 13(2):349–60. <https://doi.org/10.1590/1982-0224-20130148>
- **Casatti L.** Fish assemblage structure in a first order stream, southeastern Brazil: longitudinal distribution, seasonality, and microhabitat diversity. *Biota Neotrop*. 2005; 5(1):1–09. Available from: <https://www.biotaneotropica.org.br/v5n1/pt/abstract?article+BN02505012005>
- **Cetra M, Rondineli G, Souza UP.** Resource sharing by two neotropical freshwater fish species of streams in the Cachoeira river basin (BA). *Biota Neotrop*. 2011; 11(2):87–95. Available from: www.biotaneotropica.org.br/v11n2/en/abstract?article+bn01911022011
- **Costa ID, Rocha VM.** Feeding ecology of *Serrapinnus notomelas* (Characiformes: Cheirodontinae) in small forest streams in the Machado River basin, Rondônia, Brazil. *Acta Amazon*. 2017; 47(1):19–28. <https://doi.org/10.1590/1809-4392201601944>
- **Espírito-Santo HMV, Rodrigues MA, Zuanon J.** Reproductive strategies of Amazonian stream fishes and their fine-scale use of habitat are ordered along a hydrological gradient. *Freshwater Biol*. 2013; 58(12):2494–504. <https://doi.org/10.1111/fwb.12225>
- **Hermann TW, Stewart DJ, Limburg KE, Castello L.** Unravelling the life history of Amazonian fishes through otolith microchemistry. *R Soc Open Sci*. 2016; 3(6):160206. <http://dx.doi.org/10.1098/rsos.160206>
- **Leitão RP, Zuanon J, Mouillot D, Leal CG, Hughes RM, Kaufmann PR et al.** Disentangling the pathways of land use impacts on the functional structure of fish assemblages in Amazon streams. *Ecography*. 2018; 41(1):219–32. <https://doi.org/10.1111/ecog.02845>

- **Pease AA, González-Díaz AA, Rodiles-Hernández R, Winemiller KO.** Functional diversity and trait-environment relationships of stream fish assemblages in a large tropical catchment. *Freshw Biol.* 2012; 57(5):1060–75. <https://doi.org/10.1111/j.1365-2427.2012.02768.x>
- **Sagouis A, Jabot F, Argillier C.** Taxonomic versus functional diversity metrics: How do fish communities respond to anthropogenic stressors in reservoirs? *Ecol Freshw Fish.* 2016; 26(4):621–35. <https://doi.org/10.1111/eff.12306>
- **Santos GM, Ferreira EJG.** Peixes da bacia Amazônica. In: Lowe-McConnell RH. *Estudos ecológicos de comunidades de peixes tropicais.* São Paulo: Edusp; 1999. p.345–73.
- **Soares BE, Rosa DCO, Silva NCS, Albrecht MP, Caramaschi ÉP.** Resource use by two electric fishes (Gymnotiformes) of the National Forest Saracá-Taquera, Oriximiná, Pará. *Neotrop Ichthyol.* 2017; 15(2):e160144. <http://dx.doi.org/10.1590/1982-0224-20160144>
- **Su G, Xu J, Akasaka M, Molinos JG, Matsuzaki SS.** Human impacts on functional and taxonomic homogenization of plateau fish assemblages in Yunnan, China. *Glob Ecol Conserv.* 2015; 4:470–78. <https://doi.org/10.1016/j.gecco.2015.09.002>
- **Teresa FB, Casatti L, Cianciaruso MV.** Functional differentiation between fish assemblages from forested and deforested streams. *Neotrop Ichthyol.* 2015; 13(2):361–70. <https://doi.org/10.1590/1982-0224-20130229>
- **Villéger S, Brosse S, Mouchet M, Mouillot D, Vanni MJ.** Functional ecology of fish: current approaches and future challenges. *Aquat Sci.* 2017; 79(4):783–801. <https://doi.org/10.1007/s00027-017-0546-z>
- **Vitule JRS, Agostinho AA, Azevedo-Santos VM, Daga VS, Darwall WRT, Fitzgerald DB et al.** We need better understanding about functional diversity and vulnerability of tropical freshwater fishes. *Biodivers Conserv.* 2016; 26(3):757–62. <https://doi.org/10.1007/s10531-016-1258-8>
- **Volcan MV, Fonseca AP, Robaldo RB.** Reproduction of the threatened Annual Killifish *Austrolebias nigrofasciatus* (Cypripodontiformes: Rivulidae), confined in a natural environment. *J Threat Taxa.* 2011; 3(6):1864–67. <https://doi.org/10.11609/JoTT.o2575.1864-7>
- **Winemiller KO, Fitzgerald DB, Bower LM, Pianka ER.** Functional traits, convergent evolution, and periodic tables of niches. *Ecol Lett.* 2015; 18(8):737–51. <https://doi.org/10.1111/ele.12462>
- **Zeug SC, Winemiller KO.** Ecological correlates of fish reproductive activity in floodplain rivers: a life-history-based approach. *Can J Fish Aquat Sci.* 2007; 64(10):1291–301. <https://doi.org/10.1139/F07-094>
- **Zuanon J, Carvalho LN, Sazima I.** A chamaeleon characin: The plant-clinging and colour-changing *Ammocryptocharax elegans* (Characidiinae: Crenuchidae). *Ichthyol Explor Freshw.* 2006; 17(3):225–32. Available from: https://www.researchgate.net/publication/296937345_A_chamaeleon_characin_the_plant-clinging_and_colour-changing_Ammocryptocharax_elegans_Characidiinae_Crenuchidae
- **Zuanon J, Mendonça FP, Espírito-Santo H MV, Dias MS, Galuch AV, Akama A.** Guia de peixes da reserva Ducke – Amazônia Central. Manaus: Editora INPA; 2015. Available from: https://www.researchgate.net/publication/299459227_Guia_de_Peixes_da_Reserva_Ducke

Neotropical Ichthyology



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under
Creative Commons **CC-BY 4.0**

© 2021 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Freitas PV, Montag LFA, Ilha P, Torres NR, Maia C, Deegan L, Nascimento AT, Silva KD.** Local effects of deforestation on stream fish assemblages in the Amazon-Savannah transitional area. *Neotrop Ichthyol.* 2021; 19(3):e210098. <https://doi.org/10.1590/1982-0224-2021-0098>