

TABLE S1 | Number of samples (block nets) taken from mangrove creek fishes at seven localities in the southern Colombian Pacific coast, Panama Bight mangrove eco-region (2012–2017).

Locality (Year)	Number of samples (n)	Coordinates
Iscuandé (2014–2015)	8	2.604714, -78.058899
Mulatos (2012–2015)	10	2.648929, -78.287552
Curay (2017)	2	1.9070278, -78.5326944
Trujillo (2017)	2	1.7538611, -78.5981944
Boca Grande (2017)	2	1.8075278, -78.8478889
Teheran (2017)	2	1.6806667, -78.9559722
Chontal (2017)	2	1.5499444, -78.8890278
Total	28	



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons **CC-BY 4.0**

© 2021 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Castellanos-Galindo GA, Baos RA, Zapata LA.** Mangrove-associated fish assemblages off the southern Panama Bight region (tropical eastern Pacific). *Neotrop Ichthyol.* 2021; 19(4):e210025. <https://doi.org/10.1590/1982-0224-2021-0025>

