

TABLE S3 | Pairwise fixation index F_{ST} (below) and R_{ST} (above) in *Rhinoptera bonasus* from the coast of Brazil, between collection areas Santa Catarina (SC), São Paulo (SP), Rio de Janeiro (RJ), Pernambuco (PE) and Pará (PA). Simulation pairwise F_{ST} (below) and R_{ST} (above) between ecoregions, from Amazonic ecoregion (1, Pará – PA), Northeastern ecoregion (2, Pernambuco – PE), and Southeastern ecoregion (3, Rio de Janeiro – RJ, São Paulo – SP and Santa Catarina – SC). Bold values significant at $p < 0.001$.

		SC	SP	RJ	PE	PA
Collection areas	SC	-	0.0424	0.1344	-0.0379	0.0653
	SP	0.0049	-	0.0954	0.0265	0.0457
	RJ	0.1086	0.0756	-	0.1734	0.0448
	PE	-0.0237	0.0289	0.0948	-	0.1249
	PA	0.0398	0.1536	0.2788	0.0032	-
Ecoregions		1	2	3		
	1	-	0.6666	0.2162		
	2	0.0187	-	0.1711		
	3	0.1395	0.0032	-		



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2021 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Souza BC, Cruz VP, Almeida TRA, Sales JBL, Rodrigues-Filho LFS, Vianna M, Rotundo MM, Oliveira C, Foresti F. Genetic diversity assessment for the vulnerable migratory cownose ray *Rhinoptera bonasus* (Myliobatiformes: Rhinopteridae) from the southwestern Atlantic Ocean. Neotrop Ichthyol. 2021; 19(4):e210077. <https://doi.org/10.1590/1982-0224-2021-0077>