

TABLE S4 | Species richness estimates with rarefied (\hat{S}_{700}) and extrapolated samples (\hat{S}_{doub}) from full protection (\hat{S}_{FP}), sustainable use (\hat{S}_{SU}), and outside (\hat{S}_{Out}) with confidence intervals (parenthesis).

	\hat{S}_{FP}	\hat{S}_{SU}	\hat{S}_{Out}
\hat{S}_{700}	27.8 (24.9-30.6)	41.3 (39.3-43.2)	38.7 (36.5-40.8)
\hat{S}_{doub}	30.5 (25.1-35.9)	49.5 (43.9-55.1)	43.5 (39.1-47.8)



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2022 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Cetra M, Mattox GMT, Romero PB, Escobar SH.** Protected areas and compositional diversity of fish from Serranias Costeiras of the Ribeira de Iguape River basin, Southeast Brazil. *Neotrop Ichthyol.* 2022; 20(2):e210130. <https://doi.org/10.1590/1982-0224-2021-0130>