

TABLE S2 | Genetic pairwise differentiation between *Prochilodus lineatus* samples obtained through microsatellite markers throughout the study. *Statistically significant values $p \leq 0.007$.

Group 1	Group 2	D' Jost	F_{ST}	R_{ST}
Sep_03	Jan_05	0.62*	0.09*	0.30*
Sep_03	Aug_05	0.61*	0.13*	0.08*
Sep_03	Jan_06	0.62*	0.13*	0.29*
Sep_03	Jan_09	0.60*	0.13*	0.12*
Sep_03	Sep_10	0.57*	0.08*	0.08*
Sep_03	Feb_15	0.58*	0.08*	0.29*
Jan_05	Aug_05	0.30*	0.10*	0.50*
Jan_05	Jan_06	0.05	0.01	0.01
Jan_05	Jan_09	0.30*	0.06*	0.51*
Jan_05	Sep_10	0.33*	0.14*	0.43*
Jan_05	Feb_15	0.05	0.02	0.06
Aug_05	Jan_06	0.25*	0.06*	0.48*
Aug_05	Jan_09	0.25*	0.09*	0.28*
Aug_05	Sep_10	0.45*	0.12*	0.23*
Aug_05	Feb_15	0.27*	0.05	0.50*
Jan_06	Jan_09	0.27*	0.06*	0.44*
Jan_06	Sep_10	0.31*	0.11*	0.39*
Jan_06	Feb_15	0.03	0.00	0.00
Jan_09	Sep_10	0.47*	0.14*	0.30*
Jan_09	Feb_15	0.29*	0.04	0.41*
Sep_10	Feb_15	0.37*	0.11*	0.41*



HOW TO CITE THIS ARTICLE

- Rosa IF, Oliveira DJ, Cruz VP, Ashikaga FY, Costa GO, Doretto LB, Senhorini JA, Rocha RC, Foresti FP, Oliveira C, Foresti F. Temporal genetic structure of a stock of *Prochilodus lineatus* (Characiformes: Prochilodontidae) in the Mogi-Guaçu River ecosystem, São Paulo, Brazil. Neotrop Ichthyol. 2022; 20(2):e210156. <https://doi.org/10.1590/1982-0224-2021-0156>