

TABLE S3 | Analysis of Molecular Variance (AMOVA) of different groups of *Prochilodus lineatus*, using D-loop mitochondrial marker.*Statistically significant values $p \leq 0.08$.

Source of variation	Sum of squares	Components of variance	Percentage of variation	F' Statistics
Among the populations	19.47	0.02	1.03	$F'_{ST} : 0.01$
Within populations	477.62	2.56	98.97	
Total	497.10	2.59		



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under
Creative Commons **CC-BY 4.0**

© 2022 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Rosa IF, Oliveira DJ, Cruz VP, Ashikaga FY, Costa GO, Doretto LB, Senhorini JA, Rocha RC, Foresti FP, Oliveira C, Foresti F. Temporal genetic structure of a stock of *Prochilodus lineatus* (Characiformes: Prochilodontidae) in the Mogi-Guaçu River ecosystem, São Paulo, Brazil. Neotrop Ichthyol. 2022; 20(2):e210156. <https://doi.org/10.1590/1982-0224-2021-0156>