







**A****D****B****E****C****F**

**FIGURE S5 |** Bayesian skyline plot (BSP) analysis of temporal collections: **A.** Sep\_03, **B.** Jan\_05, **C.** Aug\_05, **D.** Jan\_06, **E.** Jan\_09, and **F.** Sep\_10 of *Prochilodus lineatus* from Mogi-Guaçu River. The y-axis, population size × generation time\*; x-axis, time (indicated in thousands of years ago). \*Generation time measured in million years. Solid lines represent median estimates, and shaded areas represent the 95% HPD limits.



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under  
Creative Commons CC-BY 4.0

© 2022 The Authors.  
Diversity and Distributions Published by SBI



Official Journal of the  
Sociedade Brasileira de Ictiologia

#### HOW TO CITE THIS ARTICLE

- Rosa IF, Oliveira DJ, Cruz VP, Ashikaga FY, Costa GO, Doretto LB, Senhorini JA, Rocha RC, Foresti FP, Oliveira C, Foresti F. Temporal genetic structure of a stock of *Prochilodus lineatus* (Characiformes: Prochilodontidae) in the Mogi-Guaçu River ecosystem, São Paulo, Brazil. Neotrop Ichthyol. 2022; 20(2):e210156. <https://doi.org/10.1590/1982-0224-2021-0156>