

TABLE S1 | Mean values and standard deviation (parentheses) of limnological characteristics for the drought and wet seasons in the control area and fish farm area at the Ilha Solteira reservoir, Upper Paraná River, São Paulo, Brazil.

Characteristics	Drought		Wet	
	Fish farm	Control	Fish Farm	Control
Δ quota (m)	-3.51 (0.34)	-3.51 (0.34)	4.05 (0.85)	4.05 (0.85)
Rainfall (mm)	5.85 (4.11)	5.84 (4.11)	14.85 (9.06)	14.85 (9.06)
Temperature (°C)	27.94 (2.28)	27.90 (2.42)	27.06 (2.75)	26.87 (1.98)
Tranparency (m)	2.50 (0.50)	3.14 (0.84)	3.69 (0.40)	4.98 (0.88)
Nitrogen ($\mu\text{g.L}^{-1}$)	0.15 (0.08)	0.05 (0.03)	0.20 (0.03)	0.24 (0.12)
Phosphorus ($\mu\text{g.L}^{-1}$)	20.57 (9.58)	20.28 (4.49)	27.61 (12.98)	11.17 (8.26)



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2022 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Neto AO, Amorim RV, Delariva RL, Camargo AFM, Veríssimo-Silveira R, Ramos IP. Structure and composition of ichthyofauna associated with cage fish farming and compared to a control area after severe drought in a Neotropical reservoir. *Neotrop Ichthyol.* 2022; 20(3):e210141. <https://doi.org/10.1590/1982-0224-2021-0141>