

**TABLE S1** | Results of Bonferroni test to detect outliers from models of food items in the diet of *Trachelyopterus galeatus*, as well as of its trophic niche breadth (CD), body condition (K) and mean degree of stomach repletion (mDR).

Model	r-student	Unadjusted p-value	Bonferroni p
Allochthonous items	1.941	0.062	NA
Terrestrial invertebrates	-3.246	0.003	0.099
Terrestrial plant	-2.091	0.045	NA
Autochthonous items	-1.941	0.062	NA
Aquatic invertebrates	-1.941	0.062	NA
Detritus	1.921	0.065	NA
Fish	3.280	0.002	0.091
CD	3.280	0.002	0.091
K	2.131	0.042	NA
mDR	3.113	0.004	0.138



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons **CC-BY 4.0**

© 2023 The Authors.  
Diversity and Distributions Published by SBI



Official Journal of the  
Sociedade Brasileira de Ictiologia

#### HOW TO CITE THIS ARTICLE

- **Bianchi-Costa IC, Quirino BA, Cardozo ALP, Yofukuji KY, Aleixo MHF, Fugi R.** Water-level fluctuations lead to changes in the diet of an omnivorous fish in a floodplain. *Neotrop Ichthyol.* 2023; 21(1):e220064. <https://doi.org/10.1590/1982-0224-2022-0064>

