

## ABGD: Automatic Barcode Gap Discovery method

/\* ABGD (Automatic Barcode Gap Discovery)

Nb seq:27 \*/

Partition 1 : found 7 groups (prior maximal distance P= 0.001000)

Partition 2 : found 7 groups (prior maximal distance P= 0.001668)

Partition 3 : found 7 groups (prior maximal distance P= 0.002783)

Partition 4 : found 7 groups (prior maximal distance P= 0.004642)

Partition 5 : found 7 groups (prior maximal distance P= 0.007743)

Partition 6 : found 7 groups (prior maximal distance P= 0.012915)

Partition 7 : found 3 groups (prior maximal distance P= 0.021544)




FIGURE S2 | Number of groups obtained with ABGD method.



OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2023 The Authors.  
Diversity and Distributions Published by SBI



Official Journal of the  
Sociedade Brasileira de Ictiologia

## HOW TO CITE THIS ARTICLE

- **Shibatta OA, Souza-Shibatta L.** New species of *Rhyacoglanis* (Siluriformes: Pseudopimelodidae) from the upper rio Tocantins basin. *Neotrop Ichthyol.* 2023; 21(1):e220075. <https://doi.org/10.1590/1982-0224-2022-0075>



Official Journal of the  
Sociedade Brasileira de Ictiologia