

ASAP method

Split probability: 0.010000 sequence length: 677

Save Spart File here

9 best partitions found by ASAP (see FAQ for more details)

The lower the score, the better the partition

Nb of species	asap-score	P-val (rank)	W (rank)	Threshold dist.	Text
* 7	2.50	8.03e-04 (1)	4.91e-04 (4)	0.012491	list csv
* 7	3.50	1.82e-01 (2)	4.75e-04 (5)	0.020786	list csv
* 8	3.50	3.07e-01 (4)	5.26e-04 (3)	0.005892	list csv
3	3.50	5.39e-01 (6)	9.35e-04 (1)	0.057153	list csv
* 4	5.00	7.37e-01 (8)	5.73e-04 (2)	0.030217	list csv
* 6	5.50	5.33e-01 (5)	4.34e-04 (6)	0.025861	list csv
14	6.00	2.30e-01 (3)	1.33e-04 (9)	0.001403	list csv
9	7.00	6.89e-01 (7)	3.96e-04 (7)	0.004215	list csv
2	8.50	9.60e-01 (9)	2.79e-04 (8)	0.087565	list csv




FIGURE S3 | Number of groups obtained with ASAP method.



OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under
Creative Commons CC-BY 4.0

© 2023 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Shibatta OA, Souza-Shibatta L. New species of *Rhyacoglanis* (Siluriformes: Pseudopimelodidae) from the upper rio Tocantins basin. Neotrop Ichthyol. 2023; 21(1):e220075. <https://doi.org/10.1590/1982-0224-2022-0075>