



FIGURE S1 | Haplotype network based on COI dataset (604 bp) showing the genetic connectivity among the 11 haplotypes of *Oligosarcus acutirostris* (24 specimens), *O. hepsetus* (one specimen), and *O. macrolepis* (two specimens). Each circle represents a unique haplotype with its size proportional to haplotype frequency. Each trace corresponds to a single mutation. Black circles represent median vectors. Colors in *O. acutirostris* correspond to the sampled river basins as indicated in the legend and showed in Fig. 2A.



OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2023 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Souza V, Oliveira-Silva L, Dutra RT, Camelier P. Phylogeography of *Oligosarcus acutirostris* (Characiformes: Characidae): testing biogeographic hypotheses in the Northeastern Mata Atlântica freshwater ecoregion. Neotrop Ichthyol. 2023; 21(2):e220084. <https://doi.org/10.1590/1982-0224-2022-0084>



Official Journal of the Sociedade Brasileira de Ictiologia