



FIGURE S3 | Residue distribution and length-weight relationship for males (A and B) and females (C and D) of *Hyphessobrycon heterorhabdus* in streams of the Guamá River basin, Eastern Amazon, State of Pará, Brazil. Black line represents models $Wt = 0.0000126 \cdot L_e^{3.12}$ for males and $Wt = 0.0001350 \cdot L_e^{2.29}$ up to 20.75 mm and $Wt = 0.0000817 \cdot L_e^{2.58}$ for females.



OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2023 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Oliveira AEP, Montag LFA, Rocha RM, López-Rodríguez NC, Prudente BS. Environmental predictors of the life history of the flag tetra *Hyphessobrycon heterorhabdus* (Characiformes: Characidae) in streams of the Eastern Amazon. Neotrop Ichthyol. 2023; 21(4):e220094. <https://doi.org/10.1590/1982-0224-2022-0094>



Official Journal of the Sociedade Brasileira de Ictiologia