

TABLE S6 | Models selected by the Akaike selection criteria (AIC) as predictors of the variation in the Gonadosomatic Index (GSI) of females of *Hyphessobrycon heterorhabdus* sampled between March 2019 and January 2020 in streams of the Guamá River basin, Eastern Amazon, State of Pará, Brazil. The bold line corresponds to the best model selected by the Akaike criteria.

	Intercept	Electrical conductivity	Accumulated monthly rainfall	Structural complexity of the substrate	Dissolved Oxygen	pH	Temperature	Stream discharge	df	logLik	AICc	delta	weight
Females	5.166			-0.0561	0.0357				4	-119.56	247.8	0	0.309
	-17.54	-0.1867			0.0433		0.8883	-6.27	6	-117.76	249	1.2	0.169
	-1.654			-0.0458	0.0343		0.2358		5	-119.18	249.4	1.6	0.139
	5.835		0.0021	-0.0644	0.0269				5	-119.22	249.5	1.68	0.134
	-11.73	-0.1219			0.0403		0.592		5	-119.25	249.6	1.75	0.129
	3.605			-0.0488	0.0348	0.1986			5	-119.32	249.7	1.88	0.121



OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2023 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Oliveira AEP, Montag LFA, Rocha RM, López-Rodríguez NC, Prudente BS.** Environmental predictors of the life history of the flag tetra *Hyphessobrycon heterorhabdus* (Characiformes: Characidae) in streams of the Eastern Amazon. *Neotrop Ichthyol.* 2023; 21(4):e220094. <https://doi.org/10.1590/1982-0224-2022-0094>



Official Journal of the Sociedade Brasileira de Ictiologia