

TABLE S3 | Beta diversity (Whittaker) among sites and across seasons.

Site	Site	Dam spillway	Dam spillway	Dam spillway	Spring-fed	Spring-fed	Spring-fed	Drainage conf.	Drainage conf.
	Season	Dry hot	Rainy	Dry cold	Dry hot	Rainy	Dry cold	Dry hot	Rainy
Dam spillway	Dry hot	-							
Dam spillway	Rainy	0	-						
Dam spillway	Dry cold	0.60	0.60	-					
Spring-fed	Dry hot	0.25	0.25	0.71	-				
Spring-fed	Rainy	0.25	0.25	0.43	0.40	-			
Spring-fed	Dry cold	0.14	0.14	0.33	0.33	0.11	-		
Drainage confluence	Dry hot	0.33	0.33	0.50	0.27	0.09	0.20	-	
Drainage confluence	Rainy	0.25	0.25	0.43	0.40	0	0.11	0.09	-
Drainage confluence	Dry cold	0.25	0.25	0.43	0.40	0	0.11	0.09	0



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2024 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Córdova-Tapia F, Palomera-Hernández V, Camacho-Cervantes M.** Invasive poeciliids dominate fish community in a highly altered river: insights from a diversity study of riverbank fishes in Mexico. *Neotrop Ichthyol.* 2024; 22(1):e230042. <https://doi.org/10.1590/1982-0224-2023-0042>

