

TABLE S1 | Morphometric and meristic data of *Austrolebias vanderbergi*. 1: MNKP 3290 (n = 5), MNKP 3293 (n = 5); 2: MNKP 3290 (n = 2), MNKP 3293 (n = 6).

	<i>Austrolebias vanderbergi</i>					
	Males ¹ (n = 10)			Females ² (n = 8)		
	Min–Max	Median (IQR)	Mean (SD)	Min–Max	Median (IQR)	Mean (SD)
Standard length (mm)	56.72–79.68	72.08 (8.92)	70.3 (6.77)	49.73–67.21	54.57 (4.62)	55.47 (5.63)
Percents of standard length						
Body depth	41.39–46.69	43.66 (2.73)	43.7 (1.97)	38.76–46.85	41.78 (0.85)	42.05 (2.34)
Caudal peduncle depth	14.39–17.07	15.79 (1.07)	15.87 (0.9)	14.44–17.07	15.78 (0.5)	15.73 (0.74)
Pre-dorsal length	45.12–54.98	49.68 (1.86)	49.69 (2.64)	53.64–59.29	56.6 (3.42)	56.34 (2.12)
Pre-pelvic length	38.49–43.19	41.56 (1.37)	41.32 (1.36)	45.02–51.91	49.71 (3.69)	49.05 (2.53)
Dorsal-fin base length	40.33–47.58	43.54 (3.68)	43.89 (2.42)	32.26–36.2	33.54 (2.47)	33.93 (1.51)
Anal-fin base length	40.14–51.14	47.45 (2.73)	47.06 (3.01)	26.91–31.79	30.05 (2.09)	29.91 (1.73)
Caudal-fin length	16.58–22.24	18.74 (2.45)	19.02 (1.76)	20.82–25.76	23.45 (2.06)	23.28 (1.59)
Pectoral-fin length	17.84–22.94	19.84 (2.28)	19.81 (1.83)	19.42–24.74	23.01 (2.81)	22.45 (1.91)
Pelvic-fin length	7.66–9.54	8.76 (1.14)	8.68 (0.69)	8.18–16.07	10.78 (1.18)	11.11 (2.29)
Head length	24.02–27.93	25.9 (1.31)	26.1 (1.22)	25.12–28.12	26.71 (0.64)	26.58 (0.95)
Percents of head length						
Head depth	122.62–136.94	131.52 (6.44)	130.67 (4.53)	117.35–130.43	122.52 (8.55)	123.13 (5.41)
Head width	68.24–75.54	71.01 (3.63)	71.8 (2.65)	71.16–77.3	75.6 (2.64)	75.04 (2.08)
Snout length	21.15–30.12	26.23 (4.32)	25.79 (2.86)	18.1–27.14	21.43 (4.9)	22.27 (3.54)
Eye diameter	21.24–26.92	24.93 (2.75)	24.69 (1.88)	22.22–31.25	28.03 (3.48)	27.36 (3.17)
Meristic counts						
Longitudinal series scales	30–33	31.5 (2)	31.2 (1.14)	28–33	29.5 (1.5)	29.75 (1.67)
Transverse series scales	14–17	16 (0)	15.7 (0.95)	14–17	15.5 (2)	15.25 (1.16)
Circumpeduncular rows scales	16–19	17.5 (1)	17.5 (1.08)	16–18	16.5 (1.25)	16.75 (0.89)
Pectoral fin rays	11–12	12 (0)	11.8 (0.42)	9–12	11 (1)	10.63 (0.92)
Pelvic fin-rays	5–5	5 (0)	5 (0)	5–5	5 (0)	5 (0)
Dorsal fin-rays	23–26	25 (2.25)	24.7 (1.25)	18–24	20.5 (2.5)	20.38 (2.07)
Anal fin-rays	25–30	27.5 (1.75)	27.6 (1.78)	22–28	23 (1.75)	23.88 (2.1)
Caudal fin-rays	19–23	21 (0.75)	21.1 (1.1)	21–24	22 (1.25)	22.25 (1.04)



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2024 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Drawert HA, Ergueta C.** Redescription of *Austrolebias accorsii* (Cyprinodontiformes: Rivulidae) and description of a new species of the genus from the upper Paraguay River basin. *Neotrop Ichthyol.* 2024; 22(2):e240001. <https://doi.org/10.1590/1982-0224-2024-0001>