



FIGURE S2 | **A.** General view of sampled streams in Parque Nacional da Amazônia, **B.** Parque Nacional do Jamanxim, **C.** Floresta Nacional Saracá-Taquera, and **D.** Reserva Biológica Nascentes da Serra do Cachimbo. Photos: **A.** Guilherme Cabral; **B.** Laboratory of Ecology and Conservation (LABECO), Federal University of Pará; **C.** Lidia Brasil Seabra; **D.** “Monitora” Project, ICMBio.



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2024 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Casatti L, Brejão GL, Carvalho FR, Costa VS, Cruz GM, Dias-Silva K, Langeani F, Lucena MDL, Oliveira-Junior JMB, Michelan TS, Montag LFA, Oliveira BC, Seabra LB, Juen L. Stream fish assemblages in the Eastern Amazon: the role of small tetras (Characidae) in alpha diversity and taxonomic structure. *Neotrop Ichthyol.* 2024; 22(3):e240014. <https://doi.org/10.1590/1982-0224-2024-0014>