

TABLE S1 | Monthly mean values of trophic niche breadth (*Ba*) and stomach repletion index (*SRI%*) for *Copella arnoldi* and *Pyrrhulina capim* from March 2019 to February 2020, as well as the trophic overlap index (Pianka) between the species.

Species	Index	2019										2020	
		Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<i>C. arnoldi</i>	<i>Ba</i>	0.67	0.60	0.58	0.31	0.56	0.49	0.60	0.40	0.56	0.56	0.60	0.77
	<i>SRI%</i>	9.87	11.37	11.17	11.10	10.75	10.78	11.70	8.85	10.54	14.12	12.95	7.11
<i>P. capim</i>	<i>Ba</i>	0.71	0.35	0.57	0.29	0.38	0.37	0.49	0.42	0.35	0.34	0.54	0.46
	<i>SRI%</i>	11.12	9.62	11.74	9.12	13.19	11.58	12.36	11.80	12.98	10.37	13.93	12.17
Trophic overlap	Pianka	0.89	0.90	0.79	0.61	0.51	0.48	0.80	0.10	0.83	0.86	0.73	0.55



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons **CC-BY 4.0**

© 2026 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Ribeiro FPT, Prudente BS, Freitas TMS.** Environmental factors drive trophic dynamics but not niche overlap of sympatric Lebiasinidae fishes in Amazonian streams. *Neotrop Ichthyol.* 2026; 24(1):e250057. <https://doi.org/10.1590/1982-0224-2025-0057>

