

FIGURE S6 | Maps colored according to the position of sub-drainage along the two Non-metric Multidimensional Scaling axes (NMDS). The NMDS have been made individually with all the corresponding dissimilarity matrices and using only two axes. In this way, each sub-drainage has a color similar to the positioning of its taxonomic, phylogenetic or trait species composition within the multivariate space.




FIGURE S6 | Continued

HOW TO CITE THIS ARTICLE

- Dias MS, Jézéquel C, Hugueny B, Leprieur F, Brosse S, Albert J, Cassemiro FAS, Frederico RG, Hidalgo M, Ortega H, Tedesco PA, Torrente-Vilara G, Zuanon J, Bigorne R, Martens K, Acosta A, Agudelo E, Maure SB, Bastos DA, Bogotá-Gregory J, Cabeceira FG, Canto ALC, Carvajal-Vallejos FM, Carvalho LN, Cella-Ribeiro A, Covain R, DoNascimento C, Dória CRC, Duarte C, Ferreira EJG, Galuch AV, Giarrizzo T, Leitão RP, Lundberg JG, Maldonado M, Montag LFA, Ohara WM, Pires THS, Pouilly M, Prada-Pedreiros S, Queiroz LJ, Py-Daniel LR, Ribeiro FRV, Herrera RR, Sarmiento J, Sousa LM, Stegmann LE, Valdiviezo-Rivera J, Villa F, Yunoki T, Wever A, Oberdorff T. Dispersal limitation, environmental filtering, and historical contingencies explain the patterns of taxonomic and phylogenetic turnover in Amazonian freshwater fish faunas, but only poorly their functional traits turnover. *Neotrop Ichthyol.* 2026; 24(1):e250084. <https://doi.org/10.1590/1982-0224-2025-0084>