

TABLE S2 | Mean and Range (%DW) of the main measurements with differences between age groups taken from adults, subadults, juveniles and neonates specimens of *Paratrygon lucindai*; (N) corresponds to the number of specimens analyzed.

<i>Paratrygon lucindai</i>	Adults			Subadults			Juveniles			Neonate	
	Mean %DW	Range %DW	N	Mean %DW	Range %DW	N	Mean %DW	Range %DW	N	Mean %DW	N
Disc width	108.8	104.5–114	24	109.2	105.4–112.1	7	109.2	103.3–113.8	16	110.2	1
Distance between snout tip and cloaca	86.4	81.2–92.2	10	85.8	82.4–88.1	5	88.4	84.9–92.3	10	92.1	1
Eye length	1.0	0.7–1.6	24	1.1	0.9–1.6	7	1.8	1.2–2.3	16	2.3	1
External length of clasper	8.1	6.4–9.8	12	6.5	5.5–7.4	2	3.0	2.6–4.0	5	3.4	1
Internal length of clasper	16.2	14.6–17.5	12	14.7	12.2–17.2	2	9.8	7.7–12.8	5	6.8	1
Pseudosiphon length	1.4	1.0–1.8	12	1.4	1.1–1.6	2	–	–	0	–	0
Ventral pseudosiphon length	6.4	5.7–7.5	12	5.7	3.0–8.4	2	–	–	0	–	0
Distance between cloaca and tail tip	37.8	24.3–81.8	21	61.6	38.7–108.3	5	88.9	37.6–141.4	14	180.2	1
Caudal sting length	7.4	4.3–10.8	14	9.0	7.4–10.3	5	11.9	9.7–14.5	9	14.7	1
Caudal sting width	0.7	0.6–1.0	19	0.7	0.6–0.9	5	0.8	0.4–1.2	11	0.6	1



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2026 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Loboda TS.** Four new species of neotropical freshwater stingrays of the genus *Paratrygon* (Myliobatiformes: Potamotrygonidae) from clear water rivers of the Amazon basin. *Neotrop Ichthyol.* 2026; 24(1):e250087. <https://doi.org/10.1590/1982-0224-2025-0087>