

TABLE S6 | Mean and Range (%DW) of the main measurements with differences between age groups taken from adults, juveniles and neonates specimens of *Paratrygon raonii*; (N) corresponds to the number of specimens analyzed.

<i>Paratrygon raonii</i>	Adults			Juveniles			Neonate		
	Mean %DW	Range %DW	N	Mean %DW	Range %DW	N	Mean %DW	Range %DW	N
Disc width	106.6	105.4–107.5	3	110.3	107–113.3	9	110.4	108.4–112.4	2
Distance between snout tip and cloaca	82	75–86.9	3	87.3	84.8–90.4	9	88.5	86–91	2
Eye length	1.4	0.9–1.6	3	2.1	1.6–2.7	9	3.1	2.8–3.4	2
External length of clasper	7.9	7.9	1	2.6	1.9–3.2	3	2.2	1.7–2.8	2
Internal length of clasper	16.5	16.5	1	8.4	7.4–10	3	7.9	7.9	2
Distance between cloaca and tail tip	40	19.5–60.5	2	87	41.5–112.2	9	65.5	35–96	2
Caudal sting length	7.7	7.7	1	11.2	8.7–15.8	8	11.1	10.3–11.9	2
Caudal sting width	0.7	0.7	1	0.8	0.7–0.9	9	1.3	1.1–1.4	2



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2026 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Loboda TS.** Four new species of neotropical freshwater stingrays of the genus *Paratrygon* (Myliobatiformes: Potamotrygonidae) from clear water rivers of the Amazon basin. *Neotrop Ichthyol.* 2026; 24(1):e250087. <https://doi.org/10.1590/1982-0224-2025-0087>

