

**TABLE S2** | List of basal production sources obtained from the Rupununi River and floodplain ponds and their functional group designations.

Water body	Basal resource	Source guild	N
Rupununi River	Filter	Seston	4
	Woody Vine	C3 Terrestrial Plants	1
	Grass	C4 Terrestrial Plants	3
	Guava Leaves	C3 Terrestrial Plants	1
	<i>Inga sp.</i>	C3 Terrestrial Plants	3
	Leaf Litter	C3 Terrestrial Plants	1
	Mimoseae Leaves	C3 Terrestrial Plants	2
	Rubiaceae (cf Diadae)	C3 Terrestrial Plants	3
Kumakapan 1	C3 Grass	C3 Terrestrial Plants	4
	Leaf Litter	C3 Terrestrial Plants	10
	Woody Vine	C3 Terrestrial Plants	1
	Roots	C3 Terrestrial Plants	3
	Riparian Tree Leaves	C3 Terrestrial Plants	7
	Leguminaceae Leaves	C3 Terrestrial Plants	2
Kumakapan 2	Cyperaceae Sedge	C3 Terrestrial Plants	2
	<i>Eichhornia sp.</i>	C3 Aquatic Plants	6
	Leaf Litter	C3 Terrestrial Plants	3
	<i>Paspalum sp.</i>	C4 Terrestrial Plants	1
	Wing Fruit	C3 Terrestrial Plants	2



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2026 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia

#### HOW TO CITE THIS ARTICLE

- Fry BL, Young CJ, Montaña CG, Winemiller LCK, Winemiller KO. Variation in basal production sources supporting fish trophic guilds in the channel and floodplain ponds of the Rupununi River, Guyana, during dry season isolation. Neotrop Ichthyol. 2026; 24(1):e250155. <https://doi.org/10.1590/1982-0224-2025-0155>