

TABLE S3 | Results of the similarity percentage analysis (SIMPER).

	Contribution (%)
<i>Acanthurus bahianus</i> x <i>A. chirurgus</i>	
Mouth width	25.61
Mouth height	16.54
Digestive tube length	16.49
Head height	12.77
Head width	10.61
<i>A. bahianus</i> x <i>A. coeruleus</i>	
Mouth width	24.76
Head height	22.47
Digestive tube length	13.30
Head width	11.63
Mouth height	11.56
<i>A. chirurgus</i> x <i>A. coeruleus</i>	
Mouth width	23.83
Head height	17.26
Digestive tube length	15.74
Mouth height	14.11
Head length	10.60



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2026 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Yukimitsu AH, Francini-Filho RB, Moura RL, Dias GM, Maganhe BL, Gibran FZ. Comparative feeding morphology and diet of SW Atlantic surgeonfishes. *Neotrop Ichthyol.* 2026; 24(1):e250166. <https://doi.org/10.1590/1982-0224-2025-0166>