

TABLE S5 | One-way ANOVA for species and jaw bones. Significant differences are shown in bold ($p < 0.05$). df = degrees of freedom; MS = Mean Square; F = F-statistic.

Jaw bones		df	MS	F	p
Maxilla	Species	2	2.603e-05	3.474	0.082
	Error	8	7.491e-06		
Premaxilla	Species	2	0.00052	25.18	<0.001
	Error	8	0.00002		
Articular	Species	2	2.920e-07	0.723	0.515
	Error	8	4.039e-07		
Dentary	Species	2	1.138e-04	10.44	0.005
	Error	8	1.089e-05		



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons **CC-BY 4.0**

© 2026 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- **Yukimitsu AH, Francini-Filho RB, Moura RL, Dias GM, Maganhe BL, Gibran FZ.** Comparative feeding morphology and diet of SW Atlantic surgeonfishes. *Neotrop Ichthyol.* 2026; 24(1):e250166. <https://doi.org/10.1590/1982-0224-2025-0166>