

TABLE S7 | Results of the similarity percentage analysis (SIMPER).

	Contribution (%)
<i>Acanthurus bahianus</i> x <i>A. chirurgus</i>	
Premaxilla	43.36
Dentary	34.10
Maxilla	18.94
Articular	3.6
<i>A. bahianus</i> x <i>A. coeruleus</i>	
Premaxilla	60.48
Dentary	20.52
Maxilla	16.18
Articular	2.82
<i>A. chirurgus</i> x <i>A. coeruleus</i>	
Premaxilla	58.42
Dentary	29.28
Maxilla	12.55
Articular	1.75



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2026 The Authors.  
Diversity and Distributions Published by SBI



Official Journal of the  
Sociedade Brasileira de Ictiologia

## HOW TO CITE THIS ARTICLE

- Yukimitsu AH, Francini-Filho RB, Moura RL, Dias GM, Maganhe BL, Gibran FZ. Comparative feeding morphology and diet of SW Atlantic surgeonfishes. Neotrop Ichthyol. 2026; 24(1):e250166. <https://doi.org/10.1590/1982-0224-2025-0166>